



SPIROTETRAMAT	GROUP	<b>23</b>	INSECTICIDE
---------------	-------	-----------	-------------

***FIFRA Section 24(c) Special Local Need Label***

**FOR DISTRIBUTION AND USE ONLY WITHIN GEORGIA**

**For the control of Asian Citrus Psyllids in citrus during bloom**

**Movento<sup>®</sup> MPC**

EPA Reg. No. 264-1065

SLN. No. GA-220004

**THIS LABEL IS VALID UNTIL December 31, 2027 UNLESS OTHERWISE AMENDED,  
WITHDRAWN, CANCELED, OR SUSPENDED.**

**ACTIVE INGREDIENT:**

Spirotetramat: cis-3-(2,5- dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl-ethyl carbonate.....14.50%

**OTHER INGREDIENTS:** .....85.50%

**TOTAL:** .....100.00%

MOVENTO MPC Contains 1.25 pounds spirotetramat per U.S. gallon (150 grams AI/liter)

**APPLICATION DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

**DIRECTIONS FOR USE**

Apply specified dosage of MOVENTO MPC as needed for control. Treatment must be made when insect populations begin to build and before a damaging population becomes established. **MOVENTO MPC must be tank-mixed with a spray adjuvant / additive having spreading and penetrating properties to maximize leaf uptake and systemicity of the active ingredient within treated plants; please contact your local Bayer CropScience representative or PCA for specific recommendations.**

**CITRUS FRUITS**

**Crops of Crop Group 10-10 Including:** Australian lime (desert, finger, and round), Brown River finger lime, Calamondin, Citron, Grapefruit, Japanese summer grapefruit, Kumquat, Lemon, Lime, Mediterranean mandarin, Mount White lime, New Guinea wild lime, Orange (sour and sweet), Pummelo, Russell River lime, Satsuma mandarin, Sweet lime, Tachibana orange, Tahiti lime, Tangelo, Tangerine, Tangor, Trifoliolate orange, Uniq fruit, including cultivars, varieties and/or hybrids of these commodities.

Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Asian citrus psyllid	16.0	0.16

**Foliar Application Restrictions:**

Pre-Harvest Interval (PHI): **1 day(s)**

Minimum interval between applications: **21 days**

Maximum MOVENTO MPC allowed per calendar year: **32 fl oz/A**

Maximum spirotetramat per calendar year: **0.31 lb ai/A**

For production in Georgia optimum control of target pests is obtained at application volumes up to 250 gpa; application volumes in excess of 350 gpa must be avoided.

Do not apply non-ionic surfactants in tank mix combination with MOVENTO MPC on white grapefruit.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call **1-866-99BAYER (1-866-992-2937)**.

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*



**Bayer CropScience**

**Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167  
1-866-99BAYER (1-866-992-2937)**